

PAIN & INFLAMMATION MANAGEMENT

Triamcinolone Acetonide Topical Aerosol (0.147 MG/G) *Sig: Apply 2-3 sprays to the affected area(s) 3-4 times daily as Directed.* QTY: 372 GMs
Mefenamic Acid 250mg capsules *Sig: Take two 250 mg capsules with food two times a day* QTY: 120 CAPS
Fluocinonide 0.1% Topical Cream *Sig: Apply 2-3 gms to affected area 3-4 times a daily. (1 gm = Dime Size)* QTY: 360 GMs
Pilaglis Cream *Sig: Apply 1-2 grams topically to affected area(s) up to 4 times daily. Peel off after 20 minutes* QTY: 240 GM

MUSCLE RELAXANTS & ORAL NSAID

Chlorzoxazone Tablet 250mg *Sig: Take 1 to 2 tablets by mouth 2 to 3 times a day* QTY: 180 TABS
Indomethacin 20mg capsules *Sig: Take 1-2 caps by mouth 2 to 3 times a day* QTY: 180 CAPS
Metaxalone 800 mg Tablet *Sig: Take 1 tablet by mouth 3-4 times daily as needed for muscle spasms* QTY: 120 TABS

DERMATITIS / ECZEMA / PSORIASIS/WOUND

Desoximetasone .05 Ointment *Sig: Apply 2 grams to the affected area(s) 4 times a day* QTY: 240 GM
Tacrolimus 0.03% Ointment *Sig: Apply 1 to 3 grams to affected area(s) up to 4 times daily* QTY: 360 GM
Clobetasol Propionate Lotion 0.05% *Sig: Apply 2 to 4 ML 3 to 4 times daily.* QTY: 236 ML

COUGH & ALLERGY

Benzonatate Capsules 200 mg Capsule *Sig: Take 1 capsules by mouth up to 3 times daily.* QTY: 90 CAPS

PROBIOTIC & WELLNESS

PreGenna NDC 717141-0072-30 *Sig: Take 1 tablet daily* QTY: 30 TABS

MIGRAINE & OTHER HEADACHE RXs

Dihydroergotamine mesylate QTY 4mg / ml x 8 *Sig: At first sign of migraine headache, administer 1 spray into Each nostril and lie down to rest in a dark, quiet room. In 15 minutes, another additional 1 spray can be applied to each nostril for a total maximum of 4 sprays.* QTY: 8 mL

ACNE

Adapalene 0.1% Topical Solution (NDC 49908-118-14) supplied in a 1.2g swab

QTY: 4 BOXES

Sig: 1 Swab QD twice daily for Acne Vulgaris

14 pads per box

DOSAGE AND ADMINISTRATION:

1. Remove swab from foil just before using. Adapalene Topical Solution, 0.1% should be applied once a day to affected areas.
2. Before retiring in the evening, wash and dry areas to be treated.
3. Apply a thin film of medication to the affected areas. Avoid the eyes, lips, and mucous membranes.
4. Discard swab after single use. Do not use if seal is broken.

During the early weeks of therapy, an apparent exacerbation of acne may occur. This is due to the action of the medication on previously unseen lesions and should not be considered a reason to discontinue therapy. Therapeutic results should be noticed after eight to twelve weeks of treatment.