

October 2022 Study Sheet

- [Peri-intubation Cardiovascular Collapse in Patients Who Are Critically Ill: Insights from the INTUBE Study - PubMed \(nih.gov\)](#).
Am J Respir Crit Care Med. 2022 Aug 15;206(4):449-458. doi: 10.1164/rccm.202111-2575OC.
 - Propofol was significantly associated with an increased risk of peri-intubation instability and collapse
- [Electrocardiographic findings associated with early clinical deterioration in acute pulmonary embolism - PubMed \(nih.gov\)](#) Acad Emerg Med. 2022 Jun 24. doi: 10.1111/acem.14554
 - In patients with acute PE, SVT is an independent predictor of clinical deterioration
- Managing TMJ Dislocations- 7 ways with photos!
 - [Managing Temporomandibular Joint Dislocations \(emergencymed.org.il\)](#)
- [Non-sterile gloves and dressing versus sterile gloves, dressings and drapes for suturing of traumatic wounds in the emergency department: a non-inferiority multicentre randomised controlled trial - PubMed \(nih.gov\)](#). Emerg Med J. 2022 Jul 26;emermed-2021-211540. doi: 10.1136/emered-2021-211540.
 - Unlikely to be a large infection difference between sterile gloves, dressings and drapes and non sterile gloves when suturing.

Ventra charting tip: If you are unable to get a full history, document why, but be specific (patient poor historian, dementia, severe COPD exacerbation, caregiver does not know child's or parent's history..ect) Do not document history limited by language (ipad or in person interpreter services are available 24/7). Do not document history limited by age if a parent or caregiver is available to give the history.